



Trends and Insights Report

Updated 02 September 2022

Purpose of report

This report comments on trends in the New Zealand COVID-19 outbreak, including cases, hospitalisations and mortality. It also comments on international COVID-19 trends and the latest scientific insights related to outbreak management. The report relies on data that may be subject to change or are incomplete. An unknown proportion of infections are not reported as cases, this proportion may differ by characteristics such as ethnic or deprivation group. Therefore, any differences in reported case rates must be interpreted with caution.

Executive Summary

Overall, all key measures of the outbreak are declining and are tracking lower than, for the first time, since March 2022. Reported case rates have continued to decrease nationally. Wastewater levels also continued to decline.

As of last week, hospitalisations and mortality have also continued decreasing nationally. This trend is expected to continue for the next couple weeks based on modelling and observed trends.

BA.5 is the dominant subvariant accounting for an estimated 91% of cases; this is consistent with wastewater findings. New variants BA.2.75 and BA.4.6 have been detected in the community at low levels. While we may see one of these sub-variants slowly predominate over the next few months, they are not expected to cause a distinct wave.

Over the next couple of weeks, it is probable that cases, hospitalisations and mortality will continue to decrease at a slower rate. However, as immunity decreases over time or if a substantially more transmissible variant emerges, we may expect fluctuations in case rates into the future.

Key insights

National Trends

<i>Cases</i>	The 7-day rolling average of reported case rates was 48.8 per 100,000 population for the week ending 28 August. This was a 33.6% decrease from the previous week, which was 73.6 per 100,000.
<i>Wastewater</i>	Wastewater quantification indicates a decreasing trend nationally.
<i>Hospitalisations</i>	The COVID-19 hospital admissions rate has been decreasing since the mid-July 7-day rolling average peak of 2.5 per 100,000, to a 7-day rolling average of 1.1 per 100,000 at 21 August.
<i>Mortality</i>	As of 28 August, there were 1,908 deaths attributed to COVID-19. The number of deaths attributed to COVID-19 appears to be continuing with a decreasing trend.
<i>Variants of Concern</i>	BA.5 makes up 91% and BA.4 makes up 5% of sequenced community cases.
<i>Border</i>	In the week ending 21 August, 85% of air border arrivals uploaded a RAT. Test positivity decreased to 2.5%. Prevalence of variants at the border generally reflects prevalence in the community.



Māori

<i>Cases</i>	The 7-day rolling average of reported case rates was 38.8 per 100,000 population at 28 August, lower than for European or Other, however there may be case ascertainment biases.
<i>Hospitalisations</i>	The age-standardised Māori hospitalisation rate for COVID-19 is 2.3 times higher than European or Other.
<i>Mortality</i>	The age-standardised cumulative mortality rate for Māori is 2.0 times higher than European or Other.

Pacific peoples

<i>Cases</i>	The 7-day rolling average of reported case rates was 35.1 per 100,000 population at 28 August; there is likely ascertainment bias, but also to note that this rate is not age adjusted.
<i>Hospitalisations</i>	Pacific peoples have the highest cumulative rate of hospitalisation with COVID-19 which is approximately 3.1 times higher than European or Other.
<i>Mortality</i>	Pacific peoples have the highest age-standardised cumulative mortality risk of any ethnicity, 2.6 times that of European or Other.

International Insights


Globally, in the week ending 28 August 2022, the number of weekly cases decreased by 16%, with 4.5 million new cases reported. The number of new weekly deaths decreased by 13% compared to the previous week, with over 13,500 fatalities reported.

Globally, between 29 July to 29 August 2022, 138,779 SARS-CoV-2 sequences were submitted to GISAID, with Omicron accounting for 99.6% of sequences.



Contents

Infection trends	5
Summary of evidence for infection and case ascertainment trends.....	5
Approximation of trends in underlying infection incidence.....	5
Test positivity trends among tertiary hospital admissions.....	6
Wastewater quantification.....	7
Modelled and observed trends in reported cases.....	8
Regional trends in reported cases.....	9
Reinfection.....	9
Demographic trends in reported case rates	10
Age trends over time.....	10
Ethnicity trends over time, by region and by age group.....	11
Deprivation trends over time.....	14
Trends in hospitalisation	15
Modelled and observed trends in hospital occupancy.....	15
Hospital admission trends over time.....	16
Age-standardised hospital admission risk for 01 January to 27 August 2022.....	20
Trends in mortality	23
Time trends in the number of deaths.....	23
Age-standardised mortality admission risk for 01 January to 27 August 2022.....	24
Whole Genomic Sequencing in community acquired cases	26
Whole Genomic Sequencing of community cases.....	26
Whole Genomic Sequencing of hospitalised cases.....	26
Border surveillance	27
Cases detected at the air border.....	27
Testing of border arrivals.....	28
Whole Genomic Sequencing of imported cases.....	28
International and scientific insights	30
Overseas waves and the likely impacts of new variants, policy changes, notifiable disease and waning immunity.....	30
Primary evidence on effectiveness of public health and outbreak control measures.....	32
Glossary	35
Data Sources.....	35
Data limitations.....	35



Figures

Figure 1: Regional reported weekly case rates of health care workers for weeks 17 July – 28 August 2022.....	5
Figure 2: Percent of tests positive among tertiary hospital admissions.	6
Figure 3: National weekly wastewater trends in SARS-CoV-2 genome copies per day	7
Figure 4: COVID-19 Modelling Aotearoa scenarios compared with reported cases nationally (BA.5 scenarios)	8
Figure 5: Regional 7-day rolling average of reported case rates from February to 28 August 2022	9
Figure 6: National reported case rates (7-day rolling average) by age for weeks 29 May - 28 August 2022.....	10
Figure 7: National 7-day rolling average of reported daily case rates by ethnicity for weeks 29 May – 28 August 2022.....	11
Figure 8: Regional reported weekly case rates by ethnicity for weeks 05 Jun – 28 August 2022.....	12
Figure 9: National ethnicity-specific reported weekly case rates by age group for weeks 05 Jun – 28 August 2022.....	13
Figure 10: National reported 7-day rolling average COVID-19 case rates by deprivation status for weeks 29 May – 28 August 2022	14
Figure 11: Reported case rates (per 100,000) with focus on those aged over 60 years, 13 February to 28 August 2022	15
Figure 12: COVID-19 Modelling Aotearoa hospital occupancy compared with observed occupancy	16
Figure 13: COVID-19 hospital admissions rate, 13 February to 21 August 2022	17
Figure 14: Hospital admission rates by age group, 13 February to 21 August 2022	18
Figure 15: Hospital admission rates by age and ethnicity, 21 May to 21 August 2022	19
Figure 16: Hospital admission rates by age and deprivation, 21 May to 21 August 2022	20
Figure 17: Age-standardised cumulative incidence (and 95% confidence intervals) of non-incident hospitalisation with COVID-19 by ethnicity, 01 January 2022 to 27 August 2022	21
Figure 18: Age-standardised cumulative incidence (and 95% confidence intervals) of non-incident hospitalisation with COVID-19 by deprivation, 01 January 2022 to 27 August 2022.....	22
Figure 19: Weekly death counts by cause of death, 29 May to 28 August 2022.....	23
Figure 20: Age-standardised cumulative incidence (and 95% confidence intervals) of mortality attributed to COVID-19 by ethnicity, 01 January 2022 to 21 August 2022.....	24
Figure 21: Age-standardised cumulative incidence (and 95% confidence intervals) of mortality attributed to COVID-19 by deprivation, 01 January 2022 to 27 August 2022	25
Figure 22: Frequency of Variants of Concern in community cases in New Zealand	26
Figure 23: Reported cases reported in post-arrival testing by country of flight departure, 01 April – 28 August 2022..	27
Figure 24: Percentage of positive tests in border arrivals who report RATs, 06 March – 21 August 2022.....	28
Figure 25: Completion metrics for border returnee testing and WGS for arrivals, 06 March – 21 August 2022	29



Infection trends

Summary of evidence for infection and case ascertainment trends


All evidence continues to support decline in incidence in the community: reported case rates in both healthcare workers and the general population have declined in the past five weeks; inpatient COVID-19 test positivity at tertiary hospitals reduced during this period and levels of viral ribonucleic acid (RNA) in wastewater have also been declining in all regions for the past few weeks.

The weekly healthcare workforce reported case rates in the week ending 28 August (7.0 per 1,000) remain higher than the general population (4.9 per 1,000).

Approximation of trends in underlying infection incidence

For the week ending 28 August, estimates suggest that 7.0 per 1,000 (245/35,169) of healthcare workers¹ (**Figure 1**) tested positive. While the healthcare work force is not representative of the general population, they are required to routinely test and report infection, so their trends in reported cases are more likely to reflect their underlying infection incidence.

Figure 1: Regional reported weekly case rates of health care workers for weeks 17 July – 28 August 2022



Source: Éclair/Episurv, 2359hrs 28 August 2022


¹ The population has been identified based on surveillance codes used in the healthcare workforce and the presence of previous testing data in 2022. A sensitivity check was run using at least 3 tests and while these numbers reduced, the incidence estimates remained very similar.



Test positivity trends among tertiary hospital admissions

Inpatient test positivity trends for tertiary hospital admissions² is shown in **Figure 2**. Tertiary hospital admission positivity is steadily declining with a 7-day rolling average³ of 1.7% for the week ending 28 August.

Figure 2: Percent of tests positive among tertiary hospital admissions.



Source: Tertiary hospitalisation data, NCTS and EpiSurv as at 2359hrs 28 August 2022

² These are hospital admissions who had COVID at the time of admission or while in hospital. These data are from Districts with tertiary hospitals; these Districts are Auckland, Canterbury, Southern, Counties Manukau, Waikato, Capital and Coast, Waitematā, and Northland.

³ A 7-day rolling average is an average of the daily values over the past seven days inclusive. This report uses both 7-day rolling averages and weekly rates; these differ in that the 7-day rolling average rate is on a daily scale and the weekly rate is on a weekly scale – hence, weekly rates are higher as they cover a longer time period.



Wastewater quantification

Figure 3 provides an overview of wastewater results nationally. Note that wastewater levels cannot be used to predict numbers of cases reliably as yet, but does indicate trends in the underlying infection rates.

Wastewater quantification indicates a decreasing trend nationally and across most regions.

Figure 3: National weekly wastewater trends in SARS-CoV-2 genome copies per day



Source: ESR SARS-CoV-2 in wastewater update for week ending 28 August 2022




Modelled and observed trends in reported cases

Figure 4 compares the latest COVID-19 Modelling Aotearoa group’s predictive model scenarios for the number of reported cases with the observed number of cases. The white line is the median prediction and grey areas indicate the upper and lower ranges of the prediction, and the red line indicates the observed cases.

The ‘July’ scenario assumes previous infection provides greater protection against reinfection and severe disease, consistent with emerging international evidence. It also incorporates updated data and future projections of uptake of second boosters, and an earlier transition to BA.5, consistent with the timing of cases and hospitalisations in New Zealand.

A peak was projected to occur in mid-July with daily cases rising to approximately 12,000 a day; however, the observed peak in reported cases was slightly earlier and lower than the median projection. Case numbers have continued to track near the lower bound of the model prediction. For the week ending 28 August there was an average of 2,425 cases reported daily, equivalent to a rate of 48.8 per 100,000 population. This is a 33.6% decrease from the previous week, which was 73.6 per 100,000.

Figure 4: COVID-19 Modelling Aotearoa scenarios compared with reported cases nationally (BA.5 scenarios)



Sources: COVID-19 Modelling Aotearoa Branching Process Model August 2022, and Ministry of Health reported case data 30 August 2022




Regional trends in reported cases

Figure 5 shows that case rates have decreased across all regions in the past week. The 7-day rolling average rate of reported cases for the Northern region (53.5 per 100,000) decreased by 33% in the past week, Te Manawa Taki (47.1 per 100,000) decreased by 36.6%, Central region (42.5 per 100,000) decreased by 35.1% and Te Waipounamu (48.3 per 100,000) decreased by 31.7%.

All Districts experienced decreases, between 23.8% and 43.8%, in the past week. The highest rate was in the Waitematā District (60.4 per 100,000) and the lowest rate was in the Hawke’s Bay District (34.5 per 100,000).

Figure 5: Regional 7-day rolling average of reported case rates from February to 28 August 2022



Source: NCTS/EpiSurv as at 2359hrs 28 August 2022

Reinfection

Analysis and interpretation of reinfection data are being developed.

It is important to note that these data come with several significant limitations: (1) Reinfections can only be identified if the previous infection was also reported. (2) Guidance on when to test after first infection was changed on June 30 prior to which the guidance was not to test until 90 days after first infection. This is now 28 days and, consequently, early reinfections were under-reported prior to June 30. (3) Those who have already had a first infection may be less likely to test during their second infection. (4) Reinfections are possibly more likely to be mild or asymptomatic.



Demographic trends in reported case rates

Since New Zealand entered Phase 3 of the Omicron response on 24 February 2022, the majority of testing has been through rapid antigen tests (RATs) rather than polymerase chain reaction (PCR) testing. RATs are self-administered and therefore require the individual to self-report their results, which may result in under-reporting of infections and of negative test results. The level of case ascertainment has also likely reduced due to active contact tracing becoming unsustainable early in the Omicron wave. As PCR testing is now only used in specific situations, PCR testing rates and positivity data are not representative of the current testing state of New Zealand, therefore demographic differences in case ascertainment cannot be evaluated.

Many infections are unreported, and the proportion of infections diagnosed and reported ('reported cases') may differ by age, ethnic and/or deprivation group. This means any difference must be interpreted with substantial caution.


Age trends over time

Figure 6 shows the 7-day rolling average rate for reported cases by age.

Reported case rates have decreased across all age groups in the past week. Nationally, reported case rates in the 65+ age group, those most at risk of poor health outcomes after infection, decreased 30.6% from the previous week to 50.0 per 100,000 in the week ending 28 August.

Among the other age groups, rates in the past week were lowest in the <5 and 5-14 age groups (29.6 and 27.1 per 100,000, respectively). The rate for the 15-24 age group was 44.8 per 100,000. The 25-44 and 45-64 age groups had the highest rates, 59.0 and 55.6 per 100,000 respectively.

Figure 6: National reported case rates (7-day rolling average) by age for weeks 29 May - 28 August 2022



Source: NCTS/EpiSurv as at 2359hrs 28 August 2022




Ethnicity trends over time, by region and by age group

Figure 7 shows the national 7-day rolling average rate of reported cases by ethnicity.

In the past week, reported case rates decreased for all ethnicities. The largest decrease was seen in Māori (36.8%) and the smallest decrease was seen in Asian peoples (30.8%). Rates in Asian (58.7 per 100,000) and European or Other (50.4 per 100,000) ethnicities remained higher than those for Māori (38.8 per 100,000) and Pacific peoples (35.1 per 100,000). But it should be noted that these rates are not adjusted for the differing age structures in the communities.

There were two distinct groupings of case rates by ethnicity, with Asian and European or Other tracking closely to each other, and Māori and Pacific peoples tracking closely to each other. The difference between these groupings was largest during the recent peak in cases and is now converging with all ethnicities tracking more closely, to lower case rates.

Figure 7: National 7-day rolling average of reported daily case rates by ethnicity for weeks 29 May – 28 August 2022



Source: NCTS/EpiSurv as at 2359hrs 28 August 2022



Figure 8 shows the regional 7-day rolling average rate of reported cases by ethnicity.

Trends in all regions were similar: rates were highest in Asian and European or Other ethnicities, and rates decreased across all ethnicities and regions, between 22.7% and 45.8%, in the past week.

Figure 8: Regional reported weekly case rates by ethnicity for weeks 05 Jun – 28 August 2022



Source: NCTS/EpiSurv as at 2359hrs 28 August 2022




Figure 9 shows national reported weekly case rates by ethnicity and age group.

In all age groups, trends were similar across all ethnicities: all rates decreased, between 15.5% and 44.8%, except Pacific peoples aged <5 increased by 2.0%. For the week ending 28 August, rates were highest in Asian and European or Other ethnicities. The rate in Asian people aged 15-24 (62.8 per 100,000) was notably higher than all other ethnicities 15-24 years old and the highest reported rate for any age group was also among Asian people (72.3 per 100,000 in 25-44 years old). The lowest reported case rates were in Māori aged <5 (18.2 per 100,000).

In 65+ age groups, the highest rate was in European or Other ethnicities at 51.2 per 100,000; followed by Asian at 48.2 per 100,000; the second lowest was in Māori at 44.2 per 100,000 and the lowest was in Pacific peoples at 36.4 per 100,000.

As Māori and Pacific peoples have lower life expectancies than other ethnicities in Aotearoa New Zealand, they are likely to have a higher risk for COVID-19 complications at a younger age than other ethnicities.

Figure 9: National ethnicity-specific reported weekly case rates by age group for weeks 05 Jun – 28 August 2022



Source: NCTS/EpiSurv as at 2359hrs 28 August 2022




Deprivation trends over time

Figure 10 shows the 7-day rolling average for reported case rates by deprivation level (based on NZDep2018⁴). This is an area, not individual, based measure. Areas of high deprivation are ones where there is poor access to the internet, low incomes, higher number of welfare recipients, increased unemployment, single parent families and higher prevalence of people living in rented accommodation and/or in homes that are overcrowded and damp.

Case rates have been highest in the least deprived areas for the past few months, but in the past few weeks have converged to a similar rate as areas of mid-range deprivation. Rates for all deprivation levels have continued to decrease; the 7-day rolling average to 28 August was highest in areas of least and mid-range deprivation (52.1 and 51.3 per 100,000 population respectively), followed by most deprived areas (44.3 per 100,000).

Behavioural insights evidence⁵ indicates that not knowing where to report RAT results, financial issues from having to isolate, inability to take time off work and not having a place to isolate safely impact the registering of a positive test. These issues could be exacerbated in areas of higher deprivation. Thus, it could be that the trends are affected by deprivation-associated bias in case ascertainment and are not a true reflection of underlying infection rates by deprivation level; however, data to investigate this are not currently available. It is also feasible that lower reported case rates in areas of high deprivation could be partially explained by higher infection rates earlier in the year.

Figure 10: National reported 7-day rolling average COVID-19 case rates by deprivation status for weeks 29 May – 28 August 2022



Source: NCTS/EpiSurv as at 2359hrs 28 August 2022

⁴ [Contents \(otago.ac.nz\)](https://www.otago.ac.nz/contents)


⁵ Information available on request



Trends in hospitalisation

Risk of hospitalisation (and mortality) is strongly linked with increasing age; since mid-June there have been decreases in reported case rates in those aged over 60 years, after substantial increases in early July, particularly in those aged 80+ (**Figure 11**). Reported case rates in over 60s were lower in the past week than seen at any time since mid-March. Consequently, hospital occupancy and hospital admissions have continued to decrease over the past month (**Figure 12** and **Figure 13**). Preliminary analysis indicates that 56% of cases are only reported as a case on the day of their hospitalisation.

Figure 11: Reported case rates (per 100,000) with focus on those aged over 60 years, 13 February to 28 August 2022



Source: NCTS/EpiSurv as at 2359hrs 28 August 2022

Modelled and observed trends in hospital occupancy


Figure 12 compares the latest COVID-19 Modelling Aotearoa group’s scenarios for hospital occupancy with observed occupancy. The white line is the median prediction and grey areas indicate the upper and lower ranges of the prediction, and the red line indicates the observed occupancy.

The 'July' scenario assumes previous infection provides greater protection against reinfection and severe disease, consistent with emerging international evidence. It also incorporates updated data and future projections of uptake of second boosters, and an earlier transition to BA.5, consistent with the timing of cases and hospitalisations in New Zealand.

The peak was projected to occur between late July and early August with daily hospitalisations rising to approximately 800 a day. The observed hospital occupancy rate has been similar to the median prediction; in the week ending 28 August the average daily occupancy was 264, equivalent to a rate of 7.0 per 100,000 population. This was a decrease of 28.2% from the week prior.



Figure 12: COVID-19 Modelling Aotearoa hospital occupancy compared with observed occupancy



Sources: COVID-19 Modelling Aotearoa (CMA) Branching Process Model August 2022, hospital occupancy (all COVID-19 positive people admitted as inpatients) from Daily hospital questionnaire as of 28 August 2022.

Hospital admission trends over time

The occupancy measures include all people who have had COVID-19, this data does not take into account whether their admission was “For” COVID-19, e.g., due to injury. Of those admitted to hospital with COVID-19 approximately 70% were probably due to COVID-19 in the week ending 21 August.

This proportion “For” COVID-19 is likely to reduce as more accurate data is made available. Therefore, in the past month the “For” estimate is likely to be slightly overestimated than actual rates “For” COVID-19.


As seen in **Figure 13**, the COVID-19 hospital admissions rate for COVID-19⁶ has been decreasing since mid-July, to a 7-day rolling average of 1.1 per 100,000 of population at 21 August.

Hospital admission rates by age group (**Figure 14**) were highest for those who are 80 years and older (13.0 per 100,000 of population), followed by those who are 70-79 years old (3.8 per 100,000) and those who are 60-69 years old (1.5 per 100,000). Admission rates among these age groups have continued to decline, decreasing by 37.9%, 31.3% and 17.2%, respectively, compared with the previous week ending 14 August.

⁶New hospital admissions who had COVID at the time of admission or while in hospital; excluding hospitalisations that were admitted and discharged within 24hrs. These data are from districts with tertiary hospitals, the districts are Auckland, Canterbury, Southern, Counties Manukau, Waikato, Capital and Coast, Waitematā and Northland.



Figure 13: COVID-19 hospital admissions⁷ rate, 13 February to 21 August 2022




Source: NMDS/Inpatients admissions feed as of 28 August 2022 data up to 21 August 2022

⁷ Hospital admission data comes from a combination of two data sources – the inpatient admission (IP) dataset (which only includes data from hospitals in certain districts) and the National Minimum Dataset (NMDS), which is a more comprehensive dataset, however, it is only available for two or more months after discharge. The IP records are provisional and overwritten by NMDS records as soon as the NMDS records become available. Please see Glossary for further details.

We can estimate the number of hospitalisations where COVID-19 could be the reason for the hospital admission. The ‘For’ measure excludes those who are identified as incidental. Recent trends are subject to revision. Please see glossary for further caveats. Coding of reasons for admission from the National Minimum dataset suggests that around 60% of people who were hospitalised with COVID-19 were hospitalised for a reason relating to their COVID-19 infection. These data are from districts with tertiary hospitals, the districts are Auckland, Canterbury, Southern, Counties Manukau, Waikato, Capital and Coast/Hutt, Waitematā and Northland.



Figure 14: Hospital admission rates by age group, 13 February to 21 August 2022



Source: NMDS/Inpatients admissions feed as of 28 August 2022 data up to 21 August 2022

Age is strongly associated with risk of hospitalisation with COVID-19, consequently it can mask other factors.

Figure 15 shows the 7-day rolling average for hospital admission rates by age and ethnicity, per 100,000. During the late-July, early August peak in COVID-19 hospital admissions, rates in Māori were highest in all age groups. Pacific peoples were second highest for under 60s-, 60-69- and 70-79-year-olds. European or Other had second highest rates in the 80 years and over and Asian people have consistently had the lowest rates across all age groups in that period.

For those aged 80+ Māori had the highest hospitalisation rates for the week ending 21 August, 31.8 per 100,000; followed by European or Other at 13.1 per 100,000; Pacific peoples at 12.3 per 100,000; Asian had the lowest hospitalisation rates, 6.5 per 100,000.

Of people aged 70-79, Pacific peoples had the highest hospitalisation rates, 12.2 per 100,000; followed by European or Other, and Māori, at 3.7 and 3.6 per 100,000 respectively. Asian people had the lowest hospitalisation rates, 1.8 per 100,000.

For the week ending 21 August of people aged 60-69, Māori had the highest hospitalisation rates, 3.8 per 100,000; followed by European or Other, and Pacific peoples, 1.5 and 1.3 per 100,000 respectively. Asian had the lowest hospitalisation rates, 0.8 per 100,000 (**Figure 15**).

Figure 16 shows the 7-day rolling average for hospital admission rates by age and deprivation, per 100,000. Consistently those in the highest deprivation have had the highest admission rates across time in all four age groups, and those in the lowest deprivation have had the lowest rates.

For the week ending 21 August of people in the most deprived areas, the highest hospitalisation rate was in the 80 years and over age group at 19.9 per 100,000; followed by mid-range deprivation areas at 11.3 per 100,000 and least deprived areas at 11.3 per 100,000.


For the week ending 21 August of people in the 70-79 age group, those from the mid-range deprivation areas had the highest hospitalisation rate for the week at 4.5 per 100,000; followed people from the most deprived areas at 3.5 per 100,000 and least deprived at 3.3 per 100,000.



For the week ending 21 August of people in the 60-69 age group, those from the mid-range deprivation areas had the highest hospitalisation rate for the week at 1.9 per 100,000; followed by those from the highest deprivation areas at 1.7 per 100,000 and those from the least deprived areas at 0.75 per 100,000.

Please note that the transparent bands are the result of Locally Weighted Scatterplot Smoothing (Loess)⁸; this creates a smooth trend line through the data points (**Figure 15 and 16**).

Figure 15: Hospital admission rates by age and ethnicity, 21 May to 21 August 2022




Source: NMDS/Inpatients admissions feed as of 28 August 2022 data up to 21 August 2022

⁸ <http://r-statistics.co/Loess-Regression-With-R.html>



Figure 16: Hospital admission rates by age and deprivation, 21 May to 21 August 2022



Source: NMDS/Inpatients admissions feed as of 28 August 2022 data up to 21 August 2022

Age-standardised hospital admission risk for 01 January to 27 August 2022

Unadjusted and age-adjusted risk: disparities in hospitalisation risk by ethnicity and deprivation are more clearly observed after adjusting (age-standardising⁹) for differences in age demographics.

Priority populations (Pacific peoples, Māori and those living in areas of high deprivation) that are at higher risk of experiencing poor health outcomes also tend to be systematically younger on average. As older age is also a strong risk for poor outcomes, the risk by ethnicity and deprivation can be masked. Therefore, the hospitalisation risk for these communities must be adjusted for age in order to make a more accurate comparison.

⁹ An age-standardised rate is a weighted average of the age-specific rates per 100,000 persons, where the weights are the proportions of persons in the corresponding age groups of the Māori population.




Figure 17 and 18 show age-standardised rates of hospitalisation with COVID-19 by ethnicity and by deprivation, respectively, for the time period of 01 January 2022 to 27 August 2022. Rates are standardised to the Māori population age structure. Data are limited to districts with tertiary hospitals in the inpatient dataset and exclude incidental hospital admissions.

Pacific peoples had the highest age-standardised rate, 3.1 times higher than European or Other. Māori had a rate 2.3 times higher than European or Other (**Figure 17**). There was no significant difference between the hospitalisation rates of Asian compared with European or Other.

Those most deprived had the highest age-standardised rate of hospitalisation with COVID-19 (2.6 times that of the least deprived) followed by those of mid-range deprivation (1.4 times that of the least deprived) (**Figure 18**).

Figure 17: Age-standardised cumulative incidence (and 95% confidence intervals¹⁰) of non-incidental¹¹ hospitalisation with COVID-19 by ethnicity, 01 January 2022 to 27 August 2022




Source: NCTS/EpiSurv, NMDS, Inpatient Admissions dataset and CVIP population estimates, 01 January 2022 to 27 August 2022

¹⁰ Hospitalisation and Mortality data, even based on complete counts, may be affected by random variation—hence, we use confidence intervals to account for the random variation inherent in these data. A 95% confidence interval means we are 95% confident that the rate would fall within the interval if we were to measure the number of hospitalisations/deaths again under the same circumstances.

¹¹ Non-incidental hospitalisations for COVID-19 refer to covid cases who have been hospitalised while an active covid case and have identified as like to be covid related. For further detail please refer to the glossary at the end of this report.



Figure 18: Age-standardised cumulative incidence (and 95% confidence intervals) of non-incident hospitalisation with COVID-19 by deprivation, 01 January 2022 to 27 August 2022



Source: NCTS/EpiSurv, NMDS, Inpatient Admissions dataset and CVIP population estimates 01 January 2022 to 27 August 2022

PROACTIVELY RELEASED




Trends in mortality

Time trends in the number of deaths

From March 2020 to 28 August 2022, there were 2,814 deaths among people who died within 28 days of being reported as a case and/or with the cause being attributable to COVID-19 (that is an underlying or contributory cause). Not all deaths have been formally coded by cause of death¹²; of the deaths that have been formally coded by cause of death, 1,215 (48%) were determined to have COVID-19 as the main underlying cause. COVID-19 contributed to a further 693 (28%) deaths, another 609 (24%) people died of an unrelated cause. Deaths have been declining after peaking in the last week of July, when almost 150 people died with COVID-19 as their underlying or a contributing cause.

Figure 19: Weekly death counts by cause of death¹³, 29 May to 28 August 2022



Source: Ministry of Health. All deaths where someone has died within 28 days of being reported as having a positive test result for COVID-19 are reported. This approach aligns with countries such as the United Kingdom; it ensures that all cases of COVID-19 who die are formally recorded to help provide an accurate assessment of the impact of COVID-19.

¹² All of the deaths within 28 days of a positive test report are fast-tracked for clinical/mortality coding to determine whether the infection was the underlying cause of the death, contributed to the death, or was unrelated to the death. An example of an unrelated death is a car accident; an example of a COVID-19 contributing is a person who dies who also has a pre-existing health condition.

¹³ Data are lagged and will be updated, interpreting with cautions of the most recent weeks.




Age-standardised mortality admission risk for 01 January to 27 August 2022

Figure 20 and 21 show age-standardised¹⁴ cumulative rates of COVID-19 attributed mortality in the population by ethnicity and by deprivation, from 01 January 2022 to 27 August 2022.

Pacific peoples had the highest risk of any ethnicity, 2.6 times that of European or Other. Rates for Māori were 2.0 times higher than European or Other, while Asian rates were lower than European or Other (**Figure 20**).

Those most deprived had the highest rate of COVID-19 attributed mortality (2.5 times that of the least deprived) followed by those of mid-range deprivation (1.5 times that of the least deprived) (**Figure 21**).

Figure 20: Age-standardised cumulative incidence (and 95% confidence intervals) of mortality attributed to COVID-19 by ethnicity, 01 January 2022 to 21 August 2022




Source: EpiSurv, Death Documents, The Healthcare User database, Mortality Collections database and CVIP population estimates, 01 January 2022 to 27 August 2022

¹⁴ An age-standardised rate is a weighted average of the age-specific rates per 100,000 persons, where the weights are the proportions of persons in the corresponding age groups of the Māori population.



Figure 21: Age-standardised cumulative incidence (and 95% confidence intervals) of mortality attributed to COVID-19 by deprivation, 01 January 2022 to 27 August 2022



Source: EpiSurv, Death Documents, The Healthcare User database, Mortality Collections database and CVIP population estimates, 01 January 2020 to 27 August 2022

PROACTIVELY RELEASED



Whole Genomic Sequencing in community acquired cases

Whole Genomic Sequencing of community cases


As of the 24 August, Whole Genomic Sequencing (WGS) for cases will be updated bi-weekly. This week's section is up to the week ending 21 August, as at the 24 August.

Omicron is the dominant variant in New Zealand having outcompeted Delta, which made up ~70% of all cases that had undergone WGS at the start of January 2022 and decreased to less than 10% of sequenced cases by the end of January.

In the two weeks to 19 August, variants BA.5, BA.4, BA.2, BA.4.6 and BA.2.75 were detected in community samples (first detected in late May/early June). In the past week BA.4/5 was detected at all wastewater sites; these two data sources confirm that BA.4/5 variants are circulating within the wider population. A small number of variant BA.2.75 and BA.4.6 cases continue to be detected in the community with six and ten cases respectively reported in the two weeks to 19 August. It is probable that small numbers of these sub-variants are circulating in the community, but they are unlikely to have a growth advantage over BA.5.

There is high certainty that BA.5 was largely responsible for the recent surge in case numbers across the country (and internationally). **Figure 22** shows that BA.5 made up about 91% of sequenced community cases in the past week. **Figure 22** also shows the increasing frequency of BA.5 in community samples over the past few weeks. As expected, we see a (relative) growth advantage of BA.5 over other variants. BA.4 has remained steady this week, making up 5% of cases.

Figure 22: Frequency of Variants of Concern in community cases in New Zealand



Source: ESR COVID-19 Genomics Insights Report #21, EpiSurv/Microreact 0900hrs 19 August 2022

Whole Genomic Sequencing of hospitalised cases

As of 19 August, ESR received samples from and had successfully processed 145 of the 570 PCR positive hospital cases with a report date in the two weeks to 19 August 2022. Of these 145 samples **4% were BA.2, 9% were BA.4** (including BA.4.6), **BA.2.75 <1% and 86% were BA.5.**

Please refer to the border surveillance section for information on WGS of imported cases.

Please see the caveats in the Glossary at the end of this document.



Border surveillance

Cases detected at the air border

Imported cases initially increased as travel volumes increased following the first stage of border reopening in March 2022. Detected cases then remained fairly constant through May and early to mid-June before rising again in late June. New Zealand's borders re-opened to all tourists and Visa holders on the 31st of July 2022. **Detected cases are below 100 for the first time since late June – Early July.**


With the removal of pre-departure testing from 20 June, it appears that detected cases increased from most countries. The increase was consistent with expectations that pre-departure testing halves the number of infected people boarding aircraft, and with increasing Omicron BA.5 prevalence in many source countries. In the past month, 2% to 4% of recent arrivals were reporting a positive test.

Figure 23 shows the number of RAT-positive cases in arrivals since April 2022. While pre-departure tests were required (before 20 June), most cases arrived on flights from Australia followed by the Cook Islands and Fiji, and then the United States of America. Since 20 June, most cases have been detected on flights from Australia, the United States of America, the United Arab Emirates, Singapore, the Cook Islands and Fiji.

Flights from Australia include both short-haul trans-Tasman flights and long-haul flights that transit through an Australian airport. It is no longer possible to accurately track the first country in a multi-stage voyage, as arrival cards are no longer scanned and data in the New Zealand Traveller Declaration system only records countries visited in the weeks before the Declaration is filled in.

While the increase in imported cases after 20 June was rapid, it was in line with expectations from the removal of pre-departure testing. Even at the peak of this increase, the total number of cases detected at the border was much less than the number reported each day in the community.

Figure 23: Reported cases reported in post-arrival testing by country of flight departure, 01 April – 28 August 2022



All cases in recent air arrivals to 09:54 AM, Monday 29 Aug 2022
Cases counted from midnight to midnight


Source: NCTS/EpiSurv as at 2359hrs 28 August 2022



Testing of border arrivals

Figure 24 shows that the percentage of positive RATs in border arrivals who reported a test was between 2 – 6% for the period 6 March – 31 July 2022. The percentage of border arrivals returning positive RATs has decreased in recent weeks and was 2.5% (1,401 of 56,694 arrivals who uploaded a test result) for the week ending 21 August.

Figure 24: Percentage of positive tests in border arrivals who report RATs, 06 March – 21 August 2022



Sources: NCTS/EpiSurv/Éclair as at 2359hrs 28 August 2022

Whole Genomic Sequencing of imported cases

Genomic sequencing data are lagging by 1 or 2 weeks because of the time required for recent arrivals to report a positive RAT, seek a follow-up PCR and have processing completed by ESR.

Figure 25 shows the completion metrics for border returnee testing¹⁵ and WGS from 06 March to 21 August 2022. The percentage of arrivals uploading a RAT has been steadily decreasing and was 85% (56,694 of 66,724 arrivals) in the week ending 21 August.

In the week ending 21 August, 46% of border arrivals who returned a positive RAT had a follow-up PCR test. This is a decrease from the previous week, at 52%. A case can only be referred to ESR for WGS if the traveller is referred for PCR testing and the lab sends the PCR sample for sequencing. Recently, approximately half of the reported RAT positive border arrivals were having a follow-up PCR test, and less than half of those PCR samples are having WGS completed.


In the week ending 21 August, the percentage of PCR positive border arrivals with WGS complete was 45%. This figure is close to those seen from mid-April to late June; between 40%-70%. This figure is expected to rise, as more of the recent cases are processed.

¹⁵ Testing and reporting at the border are a “high-trust” model and it is not expected that there will be 100% compliance with testing amongst travellers.

Labs are notified of all positive RAT results that are known to be from recent arrivals. However, some recent arrivals may not be reporting RAT results.



Figure 25: Completion metrics for border returnee testing and WGS for arrivals, 06 March – 21 August 2022



Sources: NCTS/EpiSurv/Éclair as at 2359hrs 28 August 2022, ESR WGS 21 August 2022¹⁶

Over 95% of the genomes sequenced at the border over the two-week period, ending 21 August, were BA.4/5 variants. These cases include reports of BA.2.75 and BA.4.6 in travellers to New Zealand. As at 9:00am 22 August, ESR had received samples from 478 of the 2,372 PCR-positive border cases with a report date in the two weeks to 19 August. Of the successfully sequenced samples, **85% were BA.5, 3% were BA.2, 10% were BA.4 and 2% were BA.2.75**. These proportions are similar to those seen in the community; see the community WGS section on **page 26**. In this reporting window, 13 BA.2.75 cases and 12 BA.4.6 cases were detected at the border – these are similar to case proportions found in the community during the same reporting window, however, there tends to be more cases of BA.4.6 at the border compared to the community.

¹⁶ Please note that WGS may not be completed/uploaded yet for more recent cases



International and scientific insights

Please note, global trends in cases and deaths should be interpreted with caution as several countries have been progressively changing COVID-19 testing strategies, resulting in lower overall numbers of tests performed and consequently lower numbers of cases detected.

Overseas waves and the likely impacts of new variants, policy changes, notifiable disease and waning immunity

Global

- Globally, in the week ending 28 August 2022, the number of weekly cases decreased by 16%, with 4.5 million new cases reported.
- The number of new weekly deaths decreased by 13% compared to the previous week, with over 13,500 fatalities reported.
- At the regional level, the number of reported new weekly cases decreased across all six regions: the African Region (13%), the European Region (20%), the Region of the Americas (13%), the South-East Asia Region (16%), the Eastern Mediterranean Region (37%), and the Western Pacific Region (15%).
- The number of new weekly deaths decreased across four of the six regions: the African Region (64%), the Eastern Mediterranean Region (35%), the European Region (30%), and the Region of the Americas (9%); while it increased in the South-East Asian Region (15%) and remained stable in the Western Pacific Region (3%).
- Globally, from 29 July to 29 August 2022, 138,779 SARS-CoV-2 sequences were shared through GISAID. Among these, 138,236 sequences were the Omicron Variant of Concern (VOC), accounting for 99.6% of sequences reported globally in the past 30 days.
- The prevalence of BA.2 descendent lineages (BA.2.X) remained stable compared to the previous week ending 13 August (2.7% in both weeks). BA.2.75, an Omicron descendent lineage under monitoring, still shows a relatively low prevalence globally, but a number of countries have observed recent increasing trends.
- BA.5 Omicron descendent lineages continue to be dominant globally, with an increase in weekly prevalence from 72.4% to 78.2%.

Sources: [World Health Organisation: Weekly epidemiological update on COVID-19 - 31 August 2022](#)

Australia

- In the 14 days up to 30 August 2022, there were 761 new cases per 100,000 population. This is a decrease from the week prior (14 days up to 23 August 2022) where there were 1,067 per 100,000 population.
- All states and territories saw decreases in rates of new cases compared to the previous week.
- Cases in Aboriginal and Torres Strait Islanders continue to steadily decrease after increasing up to early August.
- As at 30 August 2022, there are 2,933 current cases in hospital with 88 in ICU. This is a decrease from when last reported (23 August 2022) where there were 3,262 hospitalised cases. The majority of these cases were in New South Wales (1,794), Queensland (319) and Victoria (324) All three states continue to steadily decrease with regards to hospitalised cases.



Sources: [Australian Government: Coronavirus \(COVID-19\) common operating picture](#)

England

- Between 19 August 2022 and 26 August 2022 in England, 24,674 people had a confirmed positive test result. This shows a decrease of 19.6% compared to the previous 7 days.
- Between 23 August 2022 and 30 August 2022 in England, there have been 755,685 tests. This shows a decrease of 17.3% compared to the previous 7 days.
- Between 19 August 2022 and 26 August 2022, there have been 419 deaths within 28 days of a positive COVID-19 test. This shows a decrease of 38.2% compared to the previous 7 days.
- In the week up until and including 29 August, there were 4,124 COVID-19-related admissions to hospital, a decrease of 14.2% compared to the week prior.
- In the week up to and including 30 August, 13,542 received a first dose of vaccine, 33,618 received a second dose and 22,903 received a booster or third dose.

Sources: [Coronavirus \(COVID-19\) Data: UK](#)

Japan

- Japan continues to be the country with the highest number of new cases with a 7-day rolling average of 177,523 as at 31 August. However, infections appear to be plateauing.
- Deaths remain relatively constant with a 7-day rolling average of 279 deaths as of 30 August, compared to the previous week, at 282 deaths.
- The National Institute of Infectious Diseases has reported the occupancy rates of hospital beds for COVID-19 patients has risen and is affecting quality of care due to temporary closures of hospital wards and infected healthcare workers.
- Mass infections at facilities for the elderly may be responsible for the increase in deaths occurring alongside a decrease in infections generally.

Sources: [Our World in Data: Japan](#)

South Korea

- Following a peak in late March 2022, South Korea experienced a decline in cases. However, from late June onwards, cases have begun to rise again.
- The current 7-day rolling average for confirmed cases as at 31 August is 93,965
- The 7-day rolling average for confirmed deaths continues to increase, and as of 30 August is 77.14 per day. While this is lower than the March peak where the 7-day rolling average for deaths reached a peak of 359 per day it is unclear when this will begin to plateau.
- Reinfections continue to be driven by the immune-evasive BA.5 Omicron subvariant.

Sources: [Our World in Data: South Korea](#)



Primary evidence on effectiveness of public health and outbreak control measures

This section outlines some of the available literature about the effectiveness of public health and outbreak control measures. It is not intended to be a systematic review of all available evidence, but to provide an overview of available evidence.

Outbreak Management

- [A investigation on concordance of testing results self collected swabs versus those done by a healthcare worker](#) found that self-collection school-aged children and adolescents, following simple instructions, demonstrated high agreement with results following collection by health care workers.
- [A behavioural study from New Zealand looking at the impact of Compliance with COVID-19 measures](#) found that it is important to look at the strength of individuals' motivation and their beliefs about the advantages and disadvantages of policy outcomes and policy measures. They found this differentiation was useful in predicting an individual's possible behavioural responses to a measure.
- [A review of Taiwan's mitigation and containment strategy](#) found that non-pharmaceutical interventions, including public masking and social distancing, coupled with early and aggressive identification, isolation, and contact tracing to inhibit local transmission were optimal policies for public health management of COVID-19 and future emerging infectious diseases.
- [A study on behavioural decisions and risk perception](#) through monitoring the flows of information from both physical contact and social communication found that maintaining focus on awareness of risk among each individual's physical contacts promotes the greatest reduction in disease spread, but only when an individual is aware of the symptoms of a non-trivial proportion of their physical contacts.
- [A commentary in the Lancet on face masks](#) suggests that mass masking would be of particular importance for the protection of essential workers who cannot stay at home. As people return to work, mass masking might help to reduce a likely increase in transmission.
- [A research article on the efficacy of non-pharmaceutical interventions for COVID-19 in Europe](#) found that the population prevention and control measures implemented by the government had an impact on the change in the reproduction rate. Furthermore, that most effective factors in individual level prevention were a reduction of mobility/mixing.
- [A survey of COVID-19 in public transportation](#) looking at the risk of transmission and the impact of mitigation measures found that social distancing, density limits, masking and improving ventilation were effective at reducing the risk of transmission. R_{eff} decreased by 20% after the introduction of targeted testing and by 18% after extension of face-mask rules, reducing R_{eff} to 0.9 and suppressing the outbreak.
- [A evidence brief on the properties of the Omicron variants and how it affects public health measures effectiveness](#) found that the effects of early isolation, adult-focused reduction of interpersonal contact, and vaccination have different sites of action in infection spread dynamics and their combination can work synergistically.
- [A Canadian wastewater research paper](#) has noted that the lack of a quantitative framework to assess and interpret the wastewater data generated has been a major hurdle in translating wastewater data into public health action.
- [An observational study](#) on the impact of contact tracing and testing on controlling COVID-19 without lockdown in Hong Kong.



Economic, Social and Health Impacts

- [A research article on COVID-19 testing and mortality outcomes](#) between countries found that countries that developed stronger COVID-19 testing capacity at early timepoints, as measured by tests administered per case identified, experienced a slower increase of deaths per capita.
- [A preprint study](#) has noted that reinfections of COVID-19 are associated with an increase of risk of all-cause mortality, hospitalisation, and adverse health outcomes.
- [A population study](#) using a surveillance dataset that records all results of SARS-CoV-2 tests in France found a positive social gradient between deprivation and the risk of testing positive for SARS-CoV-2.
- [An evaluation](#) of COVID-19 policies in 50 different countries and territories considers both pharmaceutical and non-pharmaceutical interventions and assesses a jurisdiction's success at containing COVID-19 both prior to and after vaccination.
- [Systematic review of economic evaluations of COVID-19 interventions](#)
- [A cross-sectional study comparing OECD countries](#) in evaluating economic outcomes found that non-pharmaceutical interventions effectively contained the outbreaks and had positive impacts in lowering unemployment rates.
- [An research article on the disease-economy trade-offs under different epidemic control strategies](#) found that using targeted isolation would result in the best outcome for minimising both the risk of an epidemic and the economic downturn which accompanies (an epidemic).

Modelling

- [A modelling study look at preventing a cluster from becoming a new wave](#) in settings with zero community COVID-19 cases found that individual restriction or control strategy reduces the risk of an outbreak. They can be traded off against each other, but if too many are removed there is a danger of accumulating an unsafe level of risk. This has a particular impact on increasing downstream risks with increasing international travel.
- [A modelling study looking at the impact of non-pharmaceutical interventions](#) on controlling COVID-19 outbreak without lockdowns in Hong Kong found that delays in implementing control measures had significant impact on disease transmission.
- [A mathematical modelling study](#) assessing the impact of public compliance on non-pharmaceutical interventions with a cost-effectiveness analysis.
- [A modelling study](#) points to the role of super-spreader events in the contribution of novel variant predominance from a public health perspective, the results give weight to the need to focus NPIs on preventing large super-spreader events (10 or 20 secondary infections from single infected individual).
- [A preprint study](#) on social gatherings and transmission found that small gatherings, due to their frequency, can be important contributors to transmission dynamics.



Access to COVID-19 rapid antigen test kits

A summary of findings from the EBS team equitable access to COVID-19 healthcare report, released in August. The study is based on a sample populations response to COVID-19 healthcare access survey; in the field 27 June to 4 July 2022 - before improved access to antivirals, free masks and RAT tests was announced on 14 July 2022

Responses to “Have you use a RAT in the last 30 days?”, were 67% “yes”, of these respondents 87% noted access as being easy or very easy; While 4% found it difficult to access RAT’s, this figure was higher for those in 5–6-person households and those with education up to NCEA level 2; 9% and 7% respectively.

While the majority of respondents who required a RAT found accessing them easy, 31% reported, at least one issue when attempting to access a RAT. For 18-24 age group, people with disability and those in 7+ household, the likelihood of experiencing at least one access issue increase significantly: 58%, 58% and 56%, respectively. The most common issues related to receiving and picking up RATs, with these issues being significantly more impactful to 18-24 age group.

Noting that this study was conducted with a relatively small sample size, which decreased significantly when sub-grouped.

PROACTIVELY RELEASED



Glossary

Data Sources

Community Cases

Data on community cases are sourced from a combination of the National Contact Tracing Service (NCTS) and EpiSurv (New Zealand's public health surveillance platform).

Whole Genome Sequencing (WGS)

All information on WGS is sourced from the ESR COVID-19 Genomics Insights (CGI) Report, a weekly overview of SARS-CoV-2 genomic surveillance across the country.

Prevalence Estimates

National estimates of underlying infection incidence are based on the weekly test positivity in routinely asymptotically tested populations, assuming therefore that their positivity rates are indicative of their underlying infection rates. The populations identified for these estimates using surveillance codes provided for testing data are border, emergency and healthcare work forces, as well as hospital inpatients. Inpatient estimates are also produced based on a direct data feed from Tertiary hospitals rather than identifying inpatients in the national testing database; they are therefore more accurate than the national figures.

Wastewater quantification

Wastewater quantitation is a measure of the levels of virus circulating in the community. Because infectious individuals tend to shed vastly more viral particles than non-infectious individuals (particularly later on in the infection), the wastewater quantitation results are driven largely by infectious individuals, in the first 5-6 days of their infection. Although people can shed detectable virus for some weeks that can be detected by PCR testing, these individuals are unlikely to have a large impact on the quantitation curves.

Wastewater is analysed by ESR's Kenepuru and Christchurch Laboratories.

Data limitations

Prevalence estimates based on routinely tested populations

- The groups of routine testers that have been identified (healthcare, border and emergency workers, and hospital inpatients) are not a representative sample of New Zealanders, overall, they are higher risk of COVID-19 infection than the general population.
- The identification of these groups at a national level is based on surveillance codes, which may not be completed accurately, particularly since the introduction of RAT testing.
- The national estimate is for people who have uploaded at least one test result in the week, so will be an over-estimate if negative test results are not being recorded for these groups.
- National level estimates will be masking differing trends by region.
- Tertiary hospital inpatient data, while likely to be more accurate than the national level data, still reflect a higher-risk group, and neither the estimates nor the trends are generalisable to the rest of the population.
- The identification of these groups is based on surveillance codes, which may not be completed accurately, particularly since the introduction of RAT testing.
- The population has been identified based on ever having a surveillance code related to the respective workforce and having at least 2 tests (at least one of which was negative) in 2022. A sensitivity check was run using at least 3 tests and while these numbers reduced, the incidence estimates remained very similar.



Wastewater quantification

- Approximately 1 million people in New Zealand are not connected to reticulated wastewater systems.
- Samples may be either grab or 24-hour composite samples. Greater variability is expected with grab samples.
- While a standard method is being used, virus recovery can vary from sample to sample.
- SARS-CoV-2 RNA concentrations should not be compared between wastewater catchments.
- Day-to-day variability in SARS-CoV-2 RNA concentrations especially in smaller catchment is to be expected.

Hospital admissions data

- The Ministry will begin reporting COVID-19 hospitalisations using two datasets: the inpatient admission (IP) dataset – that only includes data from hospitals in certain regions – and the National Minimum Dataset (NMDS). Both of these datasets are patient-level, so they allow demographic and vaccination breakdowns to be calculated.
- Of the two databases, the IP is the more up-to-date data source for admissions. The data provided include a preliminary assessment of hospitalisations where COVID-19 may potentially play a role in the hospitalisation, based on the health specialty associated with the hospitalisation. The IP dataset does not have national coverage; it only covers hospitals in Auckland, Canterbury, Southern, Counties Manukau, Waikato, Capital and Coast, Waitemata and Northland. The IP dataset can be incomplete and provisional; it is subject to revision as the more comprehensive and more accurate NMDS data become available. One caveat is that the IP dataset does not have a reliable discharge date field. As such, it should only be used to report on admissions, not occupancy.
- The NMDS has several advantages: It provides national coverage and is a rich source of data, including data on demographics and an evaluation of the disease conditions associated with the hospital stay (including whether the admission was incidental, i.e., not related to COVID-19). However, the NMDS is only available after a significant data lag. The time lag for hospitalisation data can vary but can be approximately 60 days or more.
- Therefore, we are using a combination of these two databases for hospitalisation: the IP records are included as a provisional tally of more recent COVID-19 hospitalisations for a collection of hospitals, and then these records are overwritten by NMDS records, as soon as the NMDS records are available
- Note that the definition used for 'hospitalisation for COVID-19' in both the IP and NMDS tends to be inclusive. For the IP provisional data, the health specialty associated with the hospitalisation is used to estimate whether the hospital stay might be related COVID-19; hospitalisations that are highly unlikely to be related to COVID-19 are ruled out, as opposed to identifying hospitalisations that are likely to be COVID-related. As NMDS data become available, the clinical codes that retrospectively evaluate the reasons for the hospital stay are used to estimate if the stay was potentially related to COVID-19. The NMDS data are more robust estimation of hospitalisations 'for' COVID-19.



- This new method of data collection for COVID-19 has several advantages over the previous method, as it provides more robust data in a timely manner, using an automated method that is less burdensome and more reliable, and provides access to more detailed data. Most importantly, the new data method provides a timely and reliable way to estimate the number of hospitalisations where COVID-19 could be the reason for the hospital stay (admissions 'for' COVID-19, with some caveats). Moving forward, the majority of the reporting on hospitalisation will use the 'for COVID' definition as described above from the new databases.
- Nonetheless, we are also still able to estimate the number of hospitalisations 'with' COVID-19, i.e., an estimate of the number of hospitalisations that are associated with a positive test within 28 days of admission. Hence, in conjunction with the new hospitalisation data, we can also estimate the proportion of the total COVID-19 hospitalisations that are 'for' versus 'with'. Previous analysis has shown that the proportion of the total COVID-19 hospitalisations that are 'for' COVID-19 is about 68%.
- In addition, the new system also allows us to estimate the rate of COVID-19 hospital admissions per case or per capita.
- However, the new data feed cannot be used to estimate the proportion of all hospitalisations nationally that are associated with COVID-19. This is because we do not know the total number of patients that currently are in hospital in New Zealand for any reason at any given time (this information exists in NMDS, but only with a lag of a couple of months). Without this denominator data, we cannot calculate the proportion of all hospitalisations are associated with COVID-19.

Mortality Data

- Mortality data is lagged as to account for death coding delays and recent trends should be interpreted with caution.